Reconciling newborn screening and a novel splice variant in *BTD* associated with partial biotinidase deficiency: a BabySeq Project case report

Jaclyn B. Murry,^{1,2} Kalotina Machini,^{1,2} Ozge Ceyhan-Birsoy,^{1,3} Amy Kritzer,⁴ Joel B. Krier,^{5,6} Matthew S. Lebo,^{1,2,5} Shawn Fayer,⁶ Casie A. Genetti,⁷ Grace E. VanNoy,⁷ Timothy W. Yu,^{5,7,8} Pankaj B. Agrawal,^{5,7,9} Richard B. Parad,^{5,10} Ingrid A. Holm,^{5,7} Amy L. McGuire,¹¹ Robert C. Green,^{5,6,12} Alan H. Beggs,^{5,7} Heidi L. Rehm,^{1,2,5,12} and The BabySeq Project Team¹³

¹Laboratory for Molecular Medicine, Cambridge, Massachusetts 02139, USA; ²Department of Pathology, Brigham & Women's Hospital, Boston, Massachusetts 02115, USA; ³Department of Pathology, Memorial Sloan Kettering Cancer Center, New York, New York 10065, USA; ⁴Division of Genetics and Genomics, Boston Children's Hospital, Massachusetts 02115, USA; ⁵Harvard Medical School, Boston, Massachusetts 02115, USA; ⁶Division of Genetics, Department of Medicine, Brigham and Women's Hospital, Boston, Massachusetts 02115, USA; ⁷Division of Genetics and Genomics, The Manton Center for Orphan Disease Research, Boston Children's Hospital, Boston, Massachusetts 02115, USA; ⁸Department of Neurology, Boston Children's Hospital, Boston, Massachusetts 02115, USA; ⁹Division of Newborn Medicine, Boston Children's Hospital, Boston, Massachusetts 02115, USA; ¹⁰Department of Pediatric Newborn Medicine, Brigham and Women's Hospital, Boston, Massachusetts 02115, USA; ¹¹Center for Medical Ethics and Health Policy, Baylor College of Medicine, Houston, Texas 77030, USA; ¹²The Broad Institute of MIT and Harvard, Cambridge, Massachusetts 02141, USA

Abstract Here, we report a newborn female infant from the well-baby cohort of the BabySeq Project who was identified with compound heterozygous *BTD* gene variants. The two identified variants included a well-established pathogenic variant (c.1612C>T, p.Arg538Cys) that causes profound biotinidase deficiency (BTD) in homozygosity. In addition, a novel splice variant (c.44+1G>A, p.?) was identified in the invariant splice donor region of intron 1, potentially predictive of loss of function. The novel variant was predicted to impact splicing of exon 1; however, given the absence of any reported pathogenic variants in exon 1 and the presence of alternative splicing with exon 1 absent in most tissues in the GTEx database, we assigned an initial classification of uncertain significance. Follow-up medical record review of state-mandated newborn screen (NBS) results revealed an initial out-of-range biotinidase activity level. Levels from a repeat NBS sample barely passed cut-off into the normal range. To determine whether the infant was biotinidase-deficient, subsequent diagnostic enzyme activity testing was performed, confirming partial BTD, and resulted in a change of management for this patient. This led to reclassification of the novel splice variant based on these results. In conclusion, combining the genetic and NBS results together prompted clinical follow-up that confirmed partial BTD and informed this novel splice site's reclassification, emphasizing the importance of combining iterative genetic and phenotypic evaluations.

Corresponding author:
hrehm@bwh.harvard.edu

© 2018 Murry et al. This article is distributed under the terms of the Creative Commons Attribution License, which permits unrestricted reuse and redistribution provided that the original author and source are credited.

Ontology terms: abnormal enzyme/coenzyme activity

Published by Cold Spring Harbor Laboratory Press

doi: 10.1101/mcs.a002873

[Supplemental material is available for this article.]

¹³A complete list of the BabySeq Project Team participants appears at the end of this paper.

CASE PRESENTATION

Here we describe a newborn female who was delivered via vaginal delivery at 40 wks 5 d gestational age to a 29-yr-old mother after an uncomplicated pregnancy. The physical exam at birth was unremarkable, and all growth parameters were proportionate and appropriate for her gestational age. She was enrolled in the well-baby cohort of the BabySeq Project and randomized to undergo whole-exome sequencing with analysis limited to genes strongly associated with pediatric-onset disorders. There were no major health concerns, surgeries, or hospitalizations between her well-baby nursery discharge and the time genomic results were reported at 2.5 mo of age. Both maternal and paternal families were of European ancestry, with no known consanguinity. Family history was significant for a 17-mo-old brother with severe eczema and two distant paternal relatives reported to have alopecia (one of whom had onset as a teenager).

TECHNICAL ANALYSIS AND METHODS

In the healthy proband, both *BTB* germline variants (p.Arg538Cys and c.44+1G>A) were identified through exome sequencing as part of the BabySeq Project. Testing was performed as described previously (Ceyhan-Birsoy et al. 2017). Variants were validated by Sanger sequencing using DNA extracted from peripheral blood-derived genomic DNA. Sanger sequencing was also performed on saliva-derived genomic DNA from parents to determine phasing for both variants. Mandated newborn screening (NBS) was conducted at The New England Newborn Screening Laboratory for the Massachusetts Newborn Screening Program (<https://nensp.umassmed.edu>), and biotinidase deficiency (BTB) was screened by an enzymatic activity assay conducted on samples extracted from dried blood spots.

VARIANT INTERPRETATION

Two heterozygous variants in the *BTB* gene encoding the biotinidase enzyme were identified in the subject. One was a known pathogenic missense variant (NM_000060.4: c.1612C>T, p.Arg538Cys) (ClinVar Variation ID: 1898; <https://www.ncbi.nlm.nih.gov/clinvar>) that has been previously reported in at least five compound heterozygous and two homozygous individuals with severe BTB. It is the second most common pathogenic variant in the *BTB* gene and is associated with profound BTB. Pomponio et al. (1997) concluded that individuals with this variant are reported to have undetectable biotinyl transferase activity in their serum and <10% of mean normal biotinyl-hydrolase activity. This variant is identified in the heterozygous state on 14/121,552 non-Finnish European chromosomes in the Genome Aggregation Database (gnomAD, <http://gnomad.broadinstitute.org>; dbSNP rs80338686) (Lek et al. 2016), which is consistent with a recessive carrier frequency. This missense variant meets criteria to be classified as pathogenic for BTB in an autosomal recessive manner based on allelic observations in cases and functional evidence (Pomponio et al. 1997) (ACMG/AMP criteria codes applied: PM3_Strong [upgraded based on multiple occurrences in *trans* with pathogenic variants], PS3).

This individual also harbored a novel splice variant in intron 1 of the *BTB* gene (NM_000060.4:c.44+1G>A, p.?) (ClinVar Variation ID: 370376; <https://www.ncbi.nlm.nih.gov/clinvar>). This variant, at the time of case assessment, had not been previously reported in patients with BTB and was absent from large population studies. This variant occurs in the invariant region ($\pm 1, 2$) of the splice consensus sequence and is predicted to abolish the

Table 1. Phenotypic features

Biotinidase deficiency clinical features ^a	Proband
Biotinidase activity level	Low
Other clinical features	Absent

^aThe list of other clinical features are based on the OMIM clinical synopsis (#253260; BTD).

splice site (Alamut Interactive Biosoftware) (see Supplemental Fig. 1A). However, GTEx data support that this exon is alternatively spliced with multiple transcripts across all tissues examined either missing this exon entirely or using alternate 5' junctions (<http://www.gtexportal.org>) (see Supplemental Fig. 1B). We also noted that there were no pathogenic variants that had been reported in exon 1. Based on the available data and the uncertainty regarding whether this variant would impact the function of the biologically relevant transcript and resulting biotinidase enzyme, the variant was initially classified as a variant of uncertain significance (VUS) (ACMG/AMP codes applied: PVS1_Supporting, PM2). PVS1 was downgraded to PVS1_Supporting because of the uncertain role of exon 1 in the required function of the gene. Sanger sequencing of parental samples demonstrated that the variants were in *trans*, with the c.1612C>T, p.Arg538Cys variant being inherited from the father and the c.44+1G>A variant being inherited from the mother.

To clarify the significance of the c.44+1G>A variant, the subject's NBS results for biotinidase activity were reviewed. The initial NBS sample sent at ~24 h of life revealed only 20% enzyme activity (reference levels 30%–100%). A repeat sample generated an in-range value of 32% (see Table 1) (reference levels 30%–100%) leading to no further follow-up based on standard protocols. After the exome sequencing revealed the suspicious *BTD* genotype, to clarify whether the infant might have partial BTD, subsequent diagnostic enzyme testing was performed on a serum sample, which confirmed a low enzyme activity level consistent with a diagnosis of partial BTD (1.5 nmol/min/ml serum) (reference levels 3.5–13.8 nmol/min/ml serum). It should be noted that mean biotinidase activity has been reported at levels of 3.49 ± 0.72 SD nmol/min/ml serum in obligate heterozygotes with profound deficiency alleles (Strovel et al. 2017), suggesting carrier status is insufficient to explain the reduced enzyme level. Based on the newborn's enzyme activity level, the baby was started on biotin supplementation.

On testing, biotinidase enzyme activity suggested that the c.44+1G>A variant sufficiently impacted function of the biotinidase enzyme to result in partial enzyme deficiency when in *trans* with a variant that is associated with profound deficiency in homozygosity. Incorporation of the biotinidase enzyme testing results enabled application of ACMG/AMP criteria code PS3 elevating the variant to Likely Pathogenic (see Table 2). Subsequently, additional submitted interpretations to ClinVar for this variant (SCV000485680.1) and three other splice variants at the same position (SCV000486767.1, SCV000485974.1, and SCV000485701.1) agree with this conclusion with the first entry citing Stanley et al. (2004), who described

Table 2. Genomic findings

Gene	Genomic location	HGVS cDNA	HGVS protein	Parent of origin	Variant interpretation
<i>BTD</i>	Chr 3:15686975 (GRCh37)	NM_000060.2: c.1612C>T	p.Arg538Cys	Paternal	Pathogenic
<i>BTD</i>	Chr 3:15643402 (GRCh37)	NM_000060.2: c.44+1G>A	p.?	Maternal	Likely pathogenic

that alternative splicing of *BTD* may play a role in regulating its tissue specificity and subcellular localization. However, all of these ClinVar submissions were based on heterozygous variant observations in individuals undergoing carrier screening (P Kang, Counsyl, pers. comm.). Furthermore, it has been reported that one child identified with a novel 107-kb contiguous deletion of three genes in homozygosity, including exon 1 of the *BTD* gene, the entire *HACL1* gene, and the 5' end of the adjacent *COLQ* gene exon 1, was profoundly affected for BTD (Senanayake et al. 2015). These authors reported that at the time of evaluation, the child's symptoms were apparently solely attributable to the BTD and markedly improved with biotin supplementation (Senanayake et al. 2015). The authors concluded that the deletion of exon 1 in the *BTD* gene results in the inability to synthesize the active enzyme product, as exon 1 contains the initiation site and leader signal sequence of the enzyme, further supporting a critical role for this exon (Senanayake et al. 2015), although a deletion encompassing the entire 5' region of the gene including exon 1 may have a different impact than a variant that impacts splicing of this exon.

Biallelic pathogenic defects in the *BTD* gene lead to BTD presenting with neurocutaneous features, typically diagnosed based on clinical presentation and enzymatic activity. Individuals with partial BTD may develop symptoms upon exposure to metabolic stress and can benefit from biotin supplementation. BTD is highly treatable; but if untreated, young children with profound BTD (<10% biotinidase activity) typically exhibit neurological abnormalities, including seizures, hypotonia, ataxia, developmental delay, vision problems, hearing loss, and cutaneous abnormalities (Wolf 1993). Older children and adolescents with profound BTD frequently exhibit motor limb weakness, spastic paresis, and decreased visual acuity (Wolf 1993). Once features develop, they are usually irreversible, even with vitamin supplementation. Individuals with partial BTD (10%–30% biotinidase activity) may have hypotonia, skin rash, and hair loss, particularly during times of stress (Wolf 1993). Therapy is inexpensive (oral free biotin), with little to no side effects, and may be required lifelong (Wolf 1993). However, it is difficult to anticipate whether this child with borderline enzyme activity levels would develop symptoms if untreated, particularly given that this condition can have variable expressivity even within the same family (Wolf 1993). Because children with partial deficiency may develop symptoms, especially when stressed, and given the potential benefit, and minimal effects, all patients with partial BTD are recommended to be treated with 5 mg of biotin per day as soon as the diagnosis is made (Wolf 1993). Therefore, this infant's treatment regimen of daily vitamin biotin supplementation was instituted primarily as a precaution because of the negligible risks and significant potential benefits associated with this treatment. It should be noted that it was the identification of the *BTD* variants in conjunction with NBS results that led to the diagnosis of partial BTD in the child, resulting in follow-up and reclassification of the variant, which would not normally have been reported as an incidental finding given its initial VUS classification. The combined phenotypic analysis and genotype results were necessary to effectively diagnose and treat the baby—a process now routinely recommended for certain metabolic diseases like BTD (Strovel et al. 2017), but not possible for most other disorders. This case also illustrates the importance of functional testing in helping to decipher whether variants like the novel splice variant are disease causing. This outcome highlights the complexity of interpreting novel splice variants in genes where loss of function is associated with the disease, but for which alternative splicing of multiple transcripts confounds variant interpretation.

SUMMARY

The *BTD* c.44+1G>A variant, initially classified as a VUS because of occurrence in an alternatively spliced exon, was identified in *trans* with *BTD* p.Arg538Cys, a pathogenic variant for

profound *BTD*. Review of the NBS results identified an initial out-of-range result for *BTD*, with borderline pass on a rescreen. Confirmation of low enzyme levels upon clinical follow-up in conjunction with genomic sequencing results led to a diagnosis of partial *BTD* and the baby was started on biotin supplementation. In conclusion, this BabySeq Project case demonstrates the utility of combining genotype and biochemical phenotype information to more effectively guide the interpretation of the novel *BTD* splice site variant (c.44+1G>A variant) as likely pathogenic.

THE BABYSEQ PROJECT TEAM PARTICIPANTS

Pankaj B. Agrawal,¹⁴ Alan H. Beggs,¹⁴ Wendi N. Betting,¹⁵ Ozge Ceyhan-Birsoy,¹⁶ Kurt D. Christensen,¹⁵ Dmitry Dukhovny,¹⁷ Shawn Fayer,¹⁵ Leslie A. Frankel,¹⁸ Casie A. Genetti,¹⁴ Chet Graham,¹⁶ Robert C. Green,¹⁵ Amanda M. Gutierrez,¹⁸ Maegan Harden,¹⁹ Ingrid A. Holm,¹⁴ Joel B. Krier,¹⁵ Matthew S. Lebo,¹⁶ Harvey L. Levy,¹⁴ Xingquan Lu,¹⁸ Kalotina Machini,¹⁶ Amy L. McGuire,¹⁸ Jaclyn B. Murry,¹⁶ Medha Naik,¹⁸ Tiffany Nguyen,¹⁵ Richard B. Parad,¹⁵ Hayley A. Peoples,¹⁸ Stacey Pereira,¹⁸ Devan Petersen,¹⁸ Uma Ramamurthy,¹⁸ Vivek Ramanathan,¹⁸ Heidi L. Rehm,¹⁶ Amy Roberts,¹⁴ Jill O. Robinson,¹⁸ Serguei Roumiantsev,²⁰ Talia S. Schwartz,¹⁴ Tina K. Truong,¹⁴ Grace E. VanNoy,¹⁴ Susan E. Waisbren,¹⁴ and Timothy W. Yu¹⁴

ADDITIONAL INFORMATION

Data Deposition and Access

The variant was deposited to ClinVar (<https://www.ncbi.nlm.nih.gov/clinvar/>) and can be found under accession number SCV000712499.1.

Ethics Statement

Informed consent was obtained for the proband and family members whose results are reported in this study. The study was approved by the Institutional Review Board of Partners Healthcare and Boston Children's Hospital under protocol #2014P001906 and has received nonsignificant risk status in a pre-Investigational Device Exemption submission to the Food and Drug Administration.

Acknowledgments

We thank the family for their willingness to participate in the research study.

Author Contributions

J.B.M. contributed to initial draft of the manuscript; K.M. and O.C.B. contributed to variant interpretation; A.K. and J.B.K. contributed to clinical assessment of the subject and family; M.S.L. contributed to bioinformatics; S.F., C.A.G., and G.E.V. contributed to patient recruitment and clinical disclosure; T.W.Y., P.B.A., and R.B.P. contributed to clinical assessment of the subject and family and clinical disclosure; I.A.H. contributed to clinical assessment of the

¹⁴Boston Children's Hospital, Boston, Massachusetts 02115, USA

¹⁵Brigham and Women's Hospital, Boston, Massachusetts 02115, USA

¹⁶Laboratory for Molecular Medicine, Cambridge, Massachusetts 02139, USA

¹⁷Oregon Health and Science University, Oregon 97239, USA

¹⁸Baylor College of Medicine, Houston, Texas 77030, USA

¹⁹Broad Institute, Massachusetts 02141, USA

²⁰Massachusetts General Hospital, Boston, Massachusetts 02115, USA

Competing Interest Statement

The authors have declared no competing interest.

Referees

Barry Wolf
Anonymous

Received February 16, 2018;
accepted in revised form
April 23, 2018.

subject and family and clinical disclosure and supervision; A.L.M. contributed to ELSI assessment of the subject and family; R.C.G. and A.H.B. contributed to supervision; H.L.R. contributed to variant interpretation, supervision and manuscript preparation. BabySeq Investigators contributed to process development, infrastructure deployment, and maintenance. All authors contributed to editing the manuscript and reviewing the final draft.

Funding

This study was supported by the National Institutes of Health under awards U19HD077671. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

REFERENCES

- Ceyhan-Birsoy O, Machini K, Lebo MS, Yu TW, Agrawal PB, Parad RB, Holm IA, McGuire A, Green RC, Beggs AH, et al. 2017. A curated gene list for reporting results of newborn genomic sequencing. *Genet Med* **19**: 809–818.
- Lek M, Karczewski KJ, Minikel EV, Samocha KE, Banks E, Fennell T, O'Donnell-Luria AH, Ware JS, Hill AJ, Cummings BB, et al. 2016. Analysis of protein-coding genetic variation in 60,706 humans. *Nature* **536**: 285–291.
- Pomponio RJ, Norrgard KJ, Hymes J, Reynolds TR, Buck GA, Baumgartner R, Suormala T, Wolf B. 1997. Arg538 to Cys mutation in a CpG dinucleotide of the human biotinidase gene is the second most common cause of profound biotinidase deficiency in symptomatic children. *Hum Genet* **99**: 506–512.
- Senanayake DN, Jasinge EA, Pindolia K, Wanigasinghe J, Monaghan K, Suchy SF, Wei S, Jaysena S, Wolf B. 2015. First contiguous gene deletion causing biotinidase deficiency: the enzyme deficiency in three Sri Lankan children. *Mol Genet Metab Rep* **2**: 81–84.
- Stanley CM, Hymes J, Wolf B. 2004. Identification of alternatively spliced human biotinidase mRNAs and putative localization of endogenous biotinidase. *Mol Genet Metab* **81**: 300–312.
- Strovel ET, Cowan TM, Scott AI, Wolf B. 2017. Laboratory diagnosis of biotinidase deficiency, 2017 update: a technical standard and guideline of the American College of Medical Genetics and Genomics. *Genet Med* **19**. doi: 10.1038/gim.2017.84.
- Wolf B. 1993. Biotinidase deficiency. 2000 Mar 24 [Updated 2016 Jun 9]. In *GeneReviews® [Internet]* (ed. Adam MP, Ardinger HH, Pagon RA, et al.), University of Washington, Seattle, WA. 1993–2018. Available from <https://www.ncbi.nlm.nih.gov/books/NBK1322/>.

Reconciling newborn screening and a novel splice variant in *BTBD* associated with partial biotinidase deficiency: a BabySeq Project case report

Jaclyn B. Murry, Kalotina Machini, Ozge Ceyhan-Birsoy, et al.

Cold Spring Harb Mol Case Stud 2018, **4**: a002873 originally published online May 4, 2018
Access the most recent version at doi:[10.1101/mcs.a002873](https://doi.org/10.1101/mcs.a002873)

Supplementary Material	http://molecularcasestudies.cshlp.org/content/suppl/2018/05/04/mcs.a002873.DC1
License	This article is distributed under the terms of the Creative Commons Attribution License, which permits unrestricted reuse and redistribution provided that the original author and source are credited.
Email Alerting Service	Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or click here .