Response to olaparib in a *PALB2* germline mutated prostate cancer and genetic events associated with resistance

Peter Horak,^{1,2,3} Joachim Weischenfeldt,^{4,24} Gunhild von Amsberg,⁵ Burkhard Beyer,⁶ Andreas Schütte,⁷ Sebastian Uhrig,^{3,8,9} Laura Gieldon,^{10,11,12} Barbara Klink,^{10,11,12} Lars Feuerbach,^{3,8} Daniel Hübschmann,^{13,14,15} Simon Kreutzfeldt,^{1,3} Christoph Heining,^{11,12,16,17} Sebastian Maier,¹⁸ Barbara Hutter,^{3,8} Roland Penzel,^{3,19} Matthias Schlesner,²⁰ Roland Eils,^{21,22} Guido Sauter,²³ Albrecht Stenzinger,^{3,19} Benedikt Brors,^{3,8} Evelin Schröck,^{10,11,12} Hanno Glimm,^{11,12,16,17,25} Stefan Fröhling,^{1,2,3,25} and Thorsten Schlomm^{24,25}

¹Department of Translational Medical Oncology, National Center for Tumor Diseases (NCT) Heidelberg and German Cancer Research Center (DKFZ), 69120 Heidelberg, Germany; ²DKFZ-Heidelberg Center for Personalized Oncology (HIPO), 69120 Heidelberg, Germany; ³German Cancer Consortium (DKTK), 69120 Heidelberg, Germany; ⁴Biotech Research & Innovation Centre (BRIC) and Finsen Laboratory, University of Copenhagen and Rigshospitalet, 2200 Copenhagen, Denmark; ⁵Hubertus Wald Tumorzentrum, University Cancer Center Hamburg (UCCH), 20251 Hamburg, Germany; ⁶Martini-Clinic, Prostate Cancer Center, University Medical Center Hamburg-Eppendorf, 20246 Hamburg, Germany; ⁷Department of Urology, St. Antonius-Hospital, 48599 Gronau, Germany; ⁸Division of Applied Bioinformatics, DKFZ and NCT Heidelberg, 69120 Heidelberg, Germany; ⁹Faculty of Biosciences, Heidelberg University, 69120 Heidelberg, Germany; ¹⁰Institute for Clinical Genetics, Faculty of Medicine Carl Gustav Carus, Technische Universität Dresden, 01307 Dresden, Germany; ¹¹NCT Dresden, 01307 Dresden, Germany; ¹²German Cancer Consortium (DKTK) Dresden and German Cancer Research Center (DKFZ), 69120 Heidelberg, Germany; ¹³Division of Theoretical Bioinformatics, DKFZ, 69120 Heidelberg, Germany; ¹⁴Department of Pediatric Immunology, Hematology and Oncology, Heidelberg University Hospital, 69120 Heidelberg, Germany; ¹⁵Division of Stem Cells and Cancer, DKFZ, Heidelberg, Germany and Heidelberg Institute for Stem Cell Technology and Experimental Medicine (HI-STEM gGmbH), 69120 Heidelberg, Germany; ¹⁶Department of Translational Medical Oncology, NCT Dresden, 01307 Dresden, Germany; ¹⁷University Hospital Carl Gustav Carus, Technische Universität Dresden, 01307 Dresden, Germany; ¹⁸Progether Prostate Cancer Network, 0349 Oslo, Norway; ¹⁹Institute of Pathology, Heidelberg University Hospital, 69120 Heidelberg, Germany; ²⁰Bioinformatics and Omics Data Analytics, DKFZ, 69120 Heidelberg, Germany; ²¹Health Data Science Unit, Bioquant, Medical Faculty, University of Heidelberg, 69120 Heidelberg, Germany; ²²Center for Digital Health, Berlin Institute of Health and Charité Universitätsmedizin Berlin, 10178 Berlin, Germany; ²³Institute of Pathology, University Medical Center Hamburg-Eppendorf, 20251 Hamburg, Germany; ²⁴Department of Urology, Charité-Universitätsmedizin Berlin, 10117 Berlin, Germany

Corresponding authors:
peter.horak@nct-heidelberg.de;
thorsten.schlomm@charite.de

© 2019 Horak et al. This article is distributed under the terms of the Creative Commons Attribution-NonCommercial License, which permits reuse and redistribution, except for commercial purposes, provided that the original author and source are credited.

Ontology term: prostate cancer

Published by Cold Spring Harbor Laboratory Press

doi:10.1101/mcs.a003657

²⁵Shared last authorship

Abstract Prostate cancers harboring DNA repair gene alterations are particularly sensitive to PARP inhibitor treatment. We report a case of an advanced prostate cancer patient profiled within the NCT-MASTER (Molecularly Aided Stratification for Tumor Eradication Research) precision oncology program using next-generation sequencing. Comprehensive genomic and transcriptomic analysis identified a pathogenic germline *PALB2* variant as well as a mutational signature associated with disturbed homologous recombination together with structural genomic rearrangements. A molecular tumor board identified a potential

benefit of targeted therapy and recommended PARP inhibition and platinum-based chemotherapy. Single-agent treatment with the PARP inhibitor olaparib as well as subsequent combination with platinum-based chemotherapy resulted in disease stabilization and substantial improvement of clinical symptoms. Upon progression, we performed whole-exome and RNA sequencing of a liver metastasis, which demonstrated up-regulation of several genes characteristic for the neuroendocrine prostate cancer phenotype as well as a novel translocation resulting in an in-frame, loss-of-function fusion of *RB1*. We suggest that multidimensional genomic characterization of prostate cancer patients undergoing PARP inhibitor therapy will be necessary to capture and understand predictive biomarkers of PARP inhibitor sensitivity and resistance.

[Supplemental material is available for this article.]

INTRODUCTION

With growing evidence for the clinical activity of PARP inhibitors in several cancer entities, mutations in genes involved in HR are increasingly recognized as possible biomarkers, although the molecular events responsible for the development of resistance are incompletely understood (Lord and Ashworth 2017). Advanced prostate cancer is often characterized by germline and somatic alterations of genes involved in DNA repair via homologous recombination (HR) (Pritchard et al. 2016), which have also been associated with sensitivity to poly(ADP-ribose) polymerase (PARP) inhibition in advanced disease. These data also suggest that a substantial proportion of advanced prostate cancer patients may display “BRCAness” and can benefit from PARP inhibition and/or platinum-based chemotherapy (Lord and Ashworth 2016). Data from a phase 2 study of olaparib in metastatic castration-resistant prostate cancer (CRPC) revealed that 88% of patients that had a response to olaparib harbored one or more deleterious alterations in DNA repair genes (Mateo et al. 2015). Here we report the first case of a pathogenic germline *PALB2* (partner and localizer of BRCA2) variant that confers sensitivity to PARP inhibition in a patient with prostate cancer.

Clinical Presentation

The patient was diagnosed with locally advanced prostate adenocarcinoma at the age of 43. His family history was not suggestive of a hereditary cancer predisposition syndrome. The prostate-specific antigen (PSA) level at diagnosis was 76 ng/mL. Because of massive bladder neck infiltration, the patient underwent initial transurethral resection, followed by 6 mo of neoadjuvant combined hormonal therapy with leuporelin and bicalutamide (PSA nadir 0.6 ng/mL) and radical retropubic prostatectomy. Even though the patient had a PSA response, postoperative staging and grading exposed a large tumor with nodal metastasis and positive surgical margins (pT4, pN1, cM0, R1) and a Gleason score of 5 + 4 = 9. Within a month after radical surgery, and despite ongoing hormonal therapy, the patient became castration-refractory and was treated with six cycles of docetaxel chemotherapy. Unfortunately, prostate-specific membrane antigen positron emission tomography after completion of chemotherapy demonstrated disseminated bone and lymph node metastases. Because of a PSA response during chemotherapy, four additional cycles of docetaxel were administered. Zoledronic acid was added to the treatment regimen. The patient developed fatigue and his general condition worsened because of progressive disease and accumulating side effects of chemotherapy (fatigue and muscular weakness leading to Eastern Cooperative Oncology Group [ECOG] and World Health Organization [WHO] Score 3). Following PSA and radiographic progression with hepatic metastases, therapy with abiraterone acetate and prednisolone was initiated, which could not bring about any treatment benefit.

Table 1. Variant table

Gene	Chromosome	HGVS cDNA	HGVS protein	Variant type	Predicted effect	dbSNP	Genotype	ClinVar
PALB2	16	NM_024675.3: c.509_510delGA	p.Arg170Ilefs	Frameshift deletion	Loss of function	rs515726123	Germline heterozygous	132267

The primary tumor from the radical prostatectomy was analyzed using whole-genome sequencing and made available for clinical interpretation within the NCT-MASTER (Molecularly Aided Stratification for Tumor Eradication Research) program (Horak et al. 2017). We identified a pathogenic, heterozygous germline *PALB2* mutation (Table 1) with somatic loss of heterozygosity in the tumor as well as a mutational signature suggestive of defective DNA repair via homologous recombination when analyzing the entirety of somatic SNVs (Alexandrov et al. 2013). This prompted a molecularly informed tumor board recommendation by the NCT MASTER molecular tumor board of PARP inhibitor treatment with or without platinum-based chemotherapy as well as genetic counseling. Single-agent treatment with the PARP inhibitor olaparib resulted in substantial improvement of clinical symptoms, and only moderate side effects (e.g., anemia CTCAE Grade 2) were observed. Remarkably, the patient restarted physical activities and his general condition improved (ECOG/WHO Score 1). However, staging after 6 wk revealed disease progression with 24.1% increase of target lesions. In addition, the PSA value increased during treatment with olaparib monotherapy from 15.2 µg/L to 110 µg/L, whereas LDH decreased from 471 U/L to 297 U/L. Because of the considerable clinical improvement, olaparib treatment was continued and cisplatin (30 mg/m² i.v. weekly) was added. Magnetic resonance imaging after 2 mo of olaparib/cisplatin therapy showed disease stabilization by Prostate Cancer Clinical Trials Working Group 3 (PCWG3) criteria (Fig. 1). Addition of cisplatin resulted in a mild decrease of PSA to 96.4 µg/L as well as of LDH to 278 U/L. However, worsening anemia (CTCAE Grade 3) required transfusion of red blood cells and we observed decreasing neutrophil counts (CTCAE Grade 2).

Three months thereafter, the lymphatic and hepatic metastases progressed and clinical symptoms worsened. At time of radiological progression PSA had further decreased to 50.9 µg/L, whereas LDH dramatically increased to 1501 U/L. Given the patient's young age

Figure 1. Magnetic resonance imaging of liver metastases (A) before and (B) after 2 mo of olaparib/cisplatin treatment. Two hepatic lesions were selected and their largest diameters measured in comparison according to RECIST 1.1 criteria as defined by PCWG3. Sum of longest diameters of the two target lesions was 6.22 cm before and 5.40 cm after olaparib/cisplatin treatment (13% decrease).

and his desire to continue treatment, several experimental salvage regimens were administered. However, no durable stabilization of disease could be achieved and the patient died 5 mo later.

Germline and Somatic Genomic Alterations

To identify molecular alterations that could inform experimental therapy in this patient, we used whole-genome and transcriptome sequencing data from the prostatectomy specimen that were obtained at diagnosis within the ICGC early-onset prostate cancer project (Weischenfeldt et al. 2013) as well as whole-exome and transcriptome sequencing data from a liver metastasis that was obtained upon progression on PARP Inhibitor therapy two and a half years later. Single nucleotide variants (SNVs), small insertions and deletions (indels), copy-number variations (CNVs), genomic structural variations, and gene fusions were interpreted with regard to their clinical significance and ranked according to therapeutic relevance.

Analysis of the primary tumor revealed 14 somatic nonsynonymous SNVs and two somatic indels including a known *TP53* missense variant, NM_000546.5:c.451C>T, p.(Pro151Ser) (rs28934874) in the DNA-binding domain, which has been previously described as a pathogenic mutation in prostate cancer (Supplemental Table S1; Shi et al. 2002).

We detected a germline frameshift deletion in *PALB2*, NM_024675.3:c.509_510del, p.Arg170Ilefs*14 (rs515726123) (Table 1), which has already been described as a pathogenic Polish founder mutation associated with a predisposition for breast and ovarian cancer (Dansonka-Mieszkowska et al. 2010). This variant demonstrated somatic loss of heterozygosity with a mutational frequency of 70% in the primary tumor tissue. *PALB2* is responsible for BRCA2 nuclear localization and DNA damage response (Xia et al. 2006) and serves as a molecular adaptor between BRCA1 and BRCA2 (Sy et al. 2009). Therefore, absence of *PALB2* leads to impaired recruitment of BRCA2 and RAD51 to sites of DNA damage, making it a bona fide tumor suppressor. Germline loss-of-function mutations of *PALB2* have been primarily associated with hereditary susceptibility to breast, ovarian, and pancreatic cancer (Jones et al. 2009; Antoniou et al. 2014; Southey et al. 2016), whereas homozygous loss of function leads to Fanconi anemia (Reid et al. 2007; Xia et al. 2007).

Interestingly, we detected no other genetic hallmarks of prostate cancer, such as a *TMPRSS2-ETS* fusion or *PTEN* inactivation. We also observed prominent amplification (total copy number of 6) of the *MYC* proto-oncogene on Chromosome 8q24 (Fig. 2A), a common alteration in prostate cancer associated with increased aggressiveness (Van Den Berg et al. 1995; Jenkins et al. 1997; Fromont et al. 2013), which was accompanied by increased *MYC* mRNA expression. Another amplicon on Chromosome 2p24 encompassed the *MYC* family member *MYCN*. However, *MYCN* mRNA expression was not elevated in the primary tumor, in agreement with the observation that *MYC* and *MYCN* activation are mutually exclusive events in advanced prostate cancer (Boutros et al. 2015) and *MYCN* expression is often associated with a neuroendocrine phenotype (Beltran et al. 2016). Of note, *MYCN* expression was increased 4.8-fold in the therapy-resistant metastasis as compared to the primary tumor. Comparison of the primary and metastatic samples demonstrated no apparent evolution of CNVs contributing to additional oncogenic events, whereas extensive genomic rearrangements were evident at both time points (Fig. 2A). We also performed a supervised analysis of known mutational signatures (Alexandrov et al. 2013) and observed a strong contribution of signature Alexandrov-COSMIC 3, which is associated with defective HR DNA repair, to the mutational catalog in both the primary tumor and the metastasis (Fig. 2B).

The liver metastasis harbored 69 somatic nonsynonymous SNVs and two somatic indels (Supplemental Table S1). The *TP53* mutation as well as five other SNVs and both indels identified in the primary tumor could not be detected in the metastasis, suggesting subclonal

Figure 2. (A) DNA copy-number profiles of the primary tumor (upper panel) and metastasis (lower panel). (B) Relative contributions of mutational signatures (Alexandrov-COSMIC, AC) in the primary tumor (upper panel) and metastasis (lower panel) depicting the prevalence of the AC3 signature. (C) Distribution of measured allele frequencies of all somatic synonymous and nonsynonymous SNVs in both samples.

selection and evolution of the primary tumor during therapy. However, the distribution of allele frequencies of synonymous and nonsynonymous SNVs in both samples was influenced mainly by the tumor cell content and similar between the two samples (Fig. 2C). Interestingly, the metastasis carried a novel translocation, resulting in an in-frame fusion of *RB1* exon 3 and exon 2 of *COL28A1*, encoding a collagen subtype, and was predicted to cause loss of retinoblastoma protein function (Fig. 3). We also observed discrete changes in the transcriptional profile, in particular up-regulation of several genes characteristic for the neuroendocrine prostate cancer phenotype such as *MYCN*, *AURKA*, *EZH2*, and *DNMT1* (Beltran et al. 2016).

Figure 3. Fusion between exon 3 of *RB1* and exon 2 of *COL28A1*. Upper panels show the sequencing coverage and exon structure of the *RB1* and *COL28A1* as well as the structure of the putative fusion transcript. Chromatogram of the Sanger sequencing showing the nucleotide sequence of the fusion region is shown below.

DISCUSSION

Our observations suggest that deleterious germline *PALB2* variants represent an actionable target in metastatic CRPC and support extended genomic testing in this patient population. CRPC has been reported to be a very heterogeneous entity and several molecular subtypes have recently been defined (Aggarwal et al. 2017). Among these, tumors with DNA repair alterations might be particularly amenable to precision oncology approaches as a plethora of DNA repair genes may be linked to PARP inhibitor sensitivity. Recent studies have established the occurrence of germline *PALB2* mutations in prostate cancer as well as the sensitivity of other *PALB2* deficient tumor entities to PARP inhibition (Erkko et al. 2007; Mateo et al. 2015). However, not every patient with DNA repair defects may experience clinical benefit, and the identification and detailed characterization of predictive biomarkers is an ongoing effort. A germline missense variant in the *FANCA* gene has been associated with exceptional response to platinum in neuroendocrine prostate cancer (Beltran et al. 2015). In our case, we observed an initial response to PARP inhibition combined with platinum-based therapy, which was followed by progression of an aggressive tumor resistant to further treatment and displaying newly acquired somatic genetic alterations typically associated with neuroendocrine prostate cancer, such as an inactivating *RB1* fusion and elevated *MYCN* expression (Tan et al. 2014; Beltran et al. 2016). We did not find evidence for restoration of HR proficiency leading to secondary resistance to olaparib. We also hypothesize that resistance to PARP inhibition in prostate adenocarcinoma with “BRCAness” might co-occur with the emergence of neuroendocrine transdifferentiation (Akamatsu et al. 2018).

In summary, this comprehensive genomic and transcriptomic analysis indicates that pathogenic germline mutations in *PALB2* are associated with hallmarks of defective HR, including extensive structural genomic rearrangements and enrichment of specific mutational signatures, and might predict response to PARP inhibition as well as platinum-based chemotherapy. We propose that multidimensional genomic characterization of prostate cancer patients undergoing PARP inhibitor therapy will be necessary to define the entire spectrum of predictive biomarkers.

METHODS

Next-Generation Sequencing, Bioinformatic Analysis, and Target Validation

Whole-genome, whole-exome, and RNA sequencing were performed as described previously (Weischenfeldt et al. 2013; Kordes et al. 2016). The fusion transcript was validated by Sanger sequencing. SNVs and indels were identified using previously reported bioinformatics pipelines (Weischenfeldt et al. 2013). Structural variants were called with CREST (Wang et al. 2011). CNVs were extracted from whole-exome sequencing data with CNVkit (version 0.8.3.dev0). For the extraction of CNVs from whole-genome sequencing data we used our in-house CNV calling pipeline ACEseq (version 1.0.189) (Kleinheinz et al. 2017). A supervised analysis of mutational signatures to determine the contributions of known mutational signatures to the mutational catalog of individual samples was performed as described previously (Dieter et al. 2017). RNA-sequencing reads were mapped to the GRCh37.p13 assembly of the human reference genome using STAR 2.4.1c with the following alignment parameters: `-outFilterMultimapNmax 10 -outFilterScoreMinOverLread 0.66 -outFilterMatchNmin 0 -outFilterMatchNminOverLread 0.66 -outFilterMismatchNmax 10 -outFilterMismatchNoverLmax 0.3 -clip5pNbases 0 -clip3pNbases 0 -clip3pAfterAdapterNbases 0 -clip3pAdapterSeq AGATCGGAAGAG -clip3pAdapterMMp 0.1 -seedSearchLmax 0 -seedSearchStartLmax 50 -alignIntronMin 21 -alignIntronMax 0 -alignMatesGapMax 0 -alignEndsType Local -alignSoftClipAtReferenceEnds No -chimSegmentMin 15 -chimScoreMin 1 -chimJunctionOverhangMin 15`. The expression of genes annotated in the GencodeV19 gene model was quantified with featureCounts (version 1.4.6-p2) (Liao et al. 2014). High-confidence gene fusion predictions were extracted from chimeric alignments produced by STAR using our in-house pipeline Arriba (version 0.8), which removes recurrent alignment artifacts, transcript variants also observed in normal tissue, and reads with low sequence complexity as well as events with short anchors or breakpoints in close proximity or a low number of supporting reads relative to the overall number of predicted events in a gene. Sequencing coverage metrics are provided in Supplemental Table S2.

ADDITIONAL INFORMATION

Data Deposition and Access

The raw analytical data have been deposited in the European Genome-phenome Archive (<https://www.ebi.ac.uk/ega/datasets>) under accession numbers EGAS00001000383 (primary tumor, patient ID EOPC-125) and EGAS00001003133 (metastasis). Oncological outcome data were collected via the Progether PROM's (patient reported outcome measurement) interface (www.progether.com).

Ethics Statement

Prostate tumor samples and a matched normal blood sample were obtained following written informed consent under an institutional review board-approved protocol covering all

aspects relevant to clinical cancer genome sequencing. This study was approved and conducted in accordance with the Declaration of Helsinki.

Author Contributions

P.H., S.U., L.F., B.H., H.G., S.F., and T.S. were involved in the conception and design of the study. P.H., J.W., G.v.A., Bu.B., A.S., S.U., L.F., B.K., L.G., D.H., S.K., C.H., B.H., and R.P. collected and assembled the data. All authors performed data analysis and interpretation. P.H., J.W., S.U., B.K., L.G., S.F., and T.S. wrote the manuscript, and all authors approved the final version and are accountable for all aspects of the work.

Competing Interest Statement

The authors have declared no competing interest.

Received November 7, 2018;
accepted in revised form
February 8, 2019.

Funding

This work was supported by grant H021 from the DKFZ-Heidelberg Center for Personalized Oncology.

REFERENCES

- Aggarwal RR, Feng FY, Small EJ. 2017. Emerging categories of disease in advanced prostate cancer and their therapeutic implications. *Oncology* **31**: 467–474.
- Akamatsu S, Inoue T, Ogawa O, Gleave ME. 2018. Clinical and molecular features of treatment-related neuroendocrine prostate cancer. *Int J Urol* **25**: 345–351. doi:10.1111/iju.13526
- Alexandrov LB, Nik-Zainal S, Wedge DC, Aparicio SA, Behjati S, Biankin AV, Bignell GR, Bolli N, Borg A, Børresen-Dale AL, et al. 2013. Signatures of mutational processes in human cancer. *Nature* **500**: 415–421. doi:10.1038/nature12477
- Antoniou AC, Casadei S, Heikkinen T, Barrowdale D, Pykäs K, Roberts J, Lee A, Subramanian D, De Leeneer K, Fostira F, et al. 2014. Breast-cancer risk in families with mutations in *PALB2*. *N Engl J Med* **371**: 497–506. doi:10.1056/NEJMoa1400382
- Beltran H, Eng K, Mosquera JM, Sigaras A, Romanel A, Rennert H, Kossai M, Pauli C, Faltas B, Fontugne J, et al. 2015. Whole-exome sequencing of metastatic cancer and biomarkers of treatment response. *JAMA Oncol* **1**: 466–474. doi:10.1001/jamaoncol.2015.1313
- Beltran H, Prandi D, Mosquera JM, Benelli M, Puca L, Cyrta J, Marotz C, Giannopoulou E, Chakravarthi BV, Varambally S, et al. 2016. Divergent clonal evolution of castration-resistant neuroendocrine prostate cancer. *Nat Med* **22**: 298–305. doi:10.1038/nm.4045
- Boutros PC, Fraser M, Harding NJ, de Borja R, Trudel D, Lalonde E, Meng A, Hennings-Yeomans PH, McPherson A, Sabelnykova VY, et al. 2015. Spatial genomic heterogeneity within localized, multifocal prostate cancer. *Nat Genet* **47**: 736–745. doi:10.1038/ng.3315
- Dansonka-Mieszkowska A, Kluska A, Moes J, Dabrowska M, Nowakowska D, Niwinska A, Derlatka P, Cendrowski K, Kupryjanczyk J. 2010. A novel germline *PALB2* deletion in Polish breast and ovarian cancer patients. *BMC Med Genet* **11**: 20. doi:10.1186/1471-2350-11-20
- Dieter SM, Heining C, Agaimy A, Huebschmann D, Bonekamp D, Hutter B, Ehrenberg KR, Fröhlich M, Schlesner M, Scholl C, et al. 2017. Mutant KIT as imatinib-sensitive target in metastatic sinonasal carcinoma. *Ann Oncol* **28**: 142–148. doi:10.1093/annonc/mdw446
- Erkko H, Xia B, Nikkilä J, Schleutker J, Syrjäkoski K, Mannermaa A, Kallioniemi A, Pykäs K, Karppinen SM, Rapakko K, et al. 2007. A recurrent mutation in *PALB2* in Finnish cancer families. *Nature* **446**: 316–319. doi:10.1038/nature05609
- Fromont G, Godet J, Peyret A, Irani J, Celhay O, Rozet F, Cathelineau X, Cussenot O. 2013. 8q24 amplification is associated with Myc expression and prostate cancer progression and is an independent predictor of recurrence after radical prostatectomy. *Hum Pathol* **44**: 1617–1623. doi:10.1016/j.humpath.2013.01.012
- Horak P, Klink B, Heining C, Gröschel S, Hutter B, Fröhlich M, Uhrig S, Hübschmann D, Schlesner M, Eils R, et al. 2017. Precision oncology based on omics data: the NCT Heidelberg experience. *Int J Cancer* **141**: 877–886. doi:10.1002/ijc.30828
- Jenkins RB, Qian J, Lieber MM, Bostwick DG. 1997. Detection of c-myc oncogene amplification and chromosomal anomalies in metastatic prostatic carcinoma by fluorescence in situ hybridization. *Cancer Res* **57**: 524–531.
- Jones S, Hruban RH, Kamiyama M, Borges M, Zhang X, Parsons DW, Lin JC, Palmisano E, Brune K, Jaffee EM, et al. 2009. Exomic sequencing identifies *PALB2* as a pancreatic cancer susceptibility gene. *Science* **324**: 217. doi:10.1126/science.1171202

- Kleinheinz K, Bludau I, Huebschmann D, Heinold M, Kensche P, Gu Z, Lopez C, Hummel M, Klapper W, Moeller P, et al. 2017. ACESeq—allele specific copy number estimation from whole genome sequencing. *bioRxiv* doi:10.1101/210807
- Kordes M, Röring M, Heining C, Braun S, Hutter B, Richter D, Geörg C, Scholl C, Gröschel S, Roth W, et al. 2016. Cooperation of BRAF^{F595L} and mutant HRAS in histiocytic sarcoma provides new insights into oncogenic BRAF signaling. *Leukemia* **30**: 937–946. doi:10.1038/leu.2015.319
- Liao Y, Smyth GK, Shi W. 2014. featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. *Bioinformatics* **30**: 923–930. doi:10.1093/bioinformatics/btt656
- Lord CJ, Ashworth A. 2016. BRCAness revisited. *Nat Rev Cancer* **16**: 110–120. doi:10.1038/nrc.2015.21
- Lord CJ, Ashworth A. 2017. PARP inhibitors: synthetic lethality in the clinic. *Science* **355**: 1152–1158. doi:10.1126/science.aam7344
- Mateo J, Carreira S, Sandhu S, Miranda S, Mossop H, Perez-Lopez R, Nava Rodrigues D, Robinson D, Omlin A, Tunariu N, et al. 2015. DNA-repair defects and olaparib in metastatic prostate cancer. *N Engl J Med* **373**: 1697–1708. doi:10.1056/NEJMoa1506859
- Pritchard CC, Mateo J, Walsh MF, De Sarkar N, Abida W, Beltran H, Garofalo A, Gulati R, Carreira S, Eeles R, et al. 2016. Inherited DNA-repair gene mutations in men with metastatic prostate cancer. *N Engl J Med* **375**: 443–453. doi:10.1056/NEJMoa1603144
- Reid S, Schindler D, Hanenberg H, Barker K, Hanks S, Kalb R, Neveling K, Kelly P, Seal S, Freund M, et al. 2007. Biallelic mutations in PALB2 cause Fanconi anemia subtype FA-N and predispose to childhood cancer. *Nat Genet* **39**: 162–164. doi:10.1038/ng1947
- Shi XB, Nesslinger NJ, Deitch AD, Gumerlock PH, deVere White RW. 2002. Complex functions of mutant p53 alleles from human prostate cancer. *Prostate* **51**: 59–72. doi:10.1002/pros.10072
- Southey MC, Goldgar DE, Winqvist R, Pykäs K, Couch F, Tischkowitz M, Foulkes WD, Dennis J, Michailidou K, van Rensburg EJ, et al. 2016. PALB2, CHEK2 and ATM rare variants and cancer risk: data from COGS. *J Med Genet* **53**: 800–811. doi:10.1136/jmedgenet-2016-103839
- Sy SM, Huen MS, Chen J. 2009. PALB2 is an integral component of the BRCA complex required for homologous recombination repair. *Proc Natl Acad Sci* **106**: 7155–7160. doi:10.1073/pnas.0811159106
- Tan HL, Sood A, Rahimi HA, Wang W, Gupta N, Hicks J, Mosier S, Gocke CD, Epstein JI, Netto GJ, et al. 2014. Rb loss is characteristic of prostatic small cell neuroendocrine carcinoma. *Clin Cancer Res* **20**: 890–903. doi:10.1158/1078-0432.CCR-13-1982
- Van Den Berg C, Guan XY, Von Hoff D, Jenkins R, Bittner, Griffin C, Kallioniemi O, Visakorpi, McGill, Herath J, et al. 1995. DNA sequence amplification in human prostate cancer identified by chromosome microdissection: potential prognostic implications. *Clin Cancer Res* **1**: 11–18.
- Wang J, Mullighan CG, Easton J, Roberts S, Heatley SL, Ma J, Rusch MC, Chen K, Harris CC, Ding L, et al. 2011. CREST maps somatic structural variation in cancer genomes with base-pair resolution. *Nat Methods* **8**: 652–654. doi:10.1038/nmeth.1628
- Weischenfeldt J, Simon R, Feuerbach L, Schlangen K, Weichenhan D, Minner S, Wuttig D, Warnatz HJ, Stehr H, Rausch T, et al. 2013. Integrative genomic analyses reveal an androgen-driven somatic alteration landscape in early-onset prostate cancer. *Cancer Cell* **23**: 159–170. doi:10.1016/j.ccr.2013.01.002
- Xia B, Sheng Q, Nakanishi K, Ohashi A, Wu J, Christ N, Liu X, Jasin M, Couch FJ, Livingston DM. 2006. Control of BRCA2 cellular and clinical functions by a nuclear partner, PALB2. *Mol Cell* **22**: 719–729. doi:10.1016/j.molcel.2006.05.022
- Xia B, Dorsman JC, Ameiziane N, de Vries Y, Rooimans MA, Sheng Q, Pals G, Errami A, Gluckman E, Llera J, et al. 2007. Fanconi anemia is associated with a defect in the BRCA2 partner PALB2. *Nat Genet* **39**: 159–161. doi:10.1038/ng1942

Response to olaparib in a *PALB2* germline mutated prostate cancer and genetic events associated with resistance

Peter Horak, Joachim Weischenfeldt, Gunhild von Amsberg, et al.

Cold Spring Harb Mol Case Stud 2019, **5**: a003657 originally published online March 4, 2019
Access the most recent version at doi:[10.1101/mcs.a003657](https://doi.org/10.1101/mcs.a003657)

Supplementary Material	http://molecularcasestudies.cshlp.org/content/suppl/2019/03/07/mcs.a003657.DC1
References	This article cites 31 articles, 8 of which can be accessed free at: http://molecularcasestudies.cshlp.org/content/5/2/a003657.full.html#ref-list-1
License	This article is distributed under the terms of the Creative Commons Attribution-NonCommercial License, which permits reuse and redistribution, except for commercial purposes, provided that the original author and source are credited.
Email Alerting Service	Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or click here .