TITLE PAGE

TITLE

Biallelic SEPSECS variants in two siblings with pontocerebellar hypoplasia type 2D underscore the relevance of splice-disrupting synonymous variants in disease.

AUTHORS

Swetha Ramadesikan¹, Scott Hickey^{2,3}, Emily De Los Reyes^{4,5}, Anup D. Patel^{4,5}, Samuel J. Franklin¹, Patrick Brennan¹, Erin Crist¹, Kristy Lee^{1,3}, Peter White^{1,3}, Kim L. McBride^{2,3,6}, Daniel C. Koboldt^{1,3}, and Richard K. Wilson^{1,3}

AFFILIATIONS

¹ Steve and Cindy Rasmussen Institute for Genomic Medicine, The Research Institute at Nationwide Children's Hospital, Columbus OH

² Division of Genetic and Genomic Medicine, Nationwide Children's Hospital, Columbus OH

³ Department of Pediatrics, The Ohio State University College of Medicine, Columbus OH

⁴ Division of Neurology, Nationwide Children's Hospital, Columbus OH

⁵ Department of Neurology, The Ohio State University College of Medicine, Columbus OH

⁶ Center for Cardiovascular Research, The Research Institute at Nationwide Children's Hospital, Columbus OH

Short Running Title:

Biallelic SEPSECS variants result in pontocerebellar hypoplasia type 2D

Corresponding author: Daniel C. Koboldt

ABSTRACT

Noncoding and synonymous coding variants that exert their effects via alternative splicing are increasingly recognized as an important category of disease-causing variants. In this report, we describe two siblings who presented with hypotonia, profound developmental delays, and seizures. Brain MRI in the proband at 5 years showed diffuse cerebral and cerebellar white matter volume loss. Both siblings later developed ventilator-dependent respiratory insufficiency, scoliosis and are currently nonverbal and non-ambulatory. Extensive molecular testing including oligo array and clinical exome sequencing was non-diagnostic. Research genome sequencing under an IRB-approved study protocol revealed that both affected children were compound-heterozygous for variants in the *SEPSECS* gene. One variant was an initiator codon change (c.1A>T) that disrupted protein translation, consistent with the observation that most disease-causing variants are loss-of-function changes. The other variant was a coding change (c.846G>A) that was predicted to be synonymous but had been demonstrated to disrupt mRNA splicing in a minigene assay.

SEPSECS gene encodes O-phosphoseryl-tRNA(Sec) selenium transferase, an enzyme that participates in the biosynthesis and transport of selenoproteins in the body. Variations in *SEPSECS* cause autosomal recessive pontocerebellar hypoplasia type 2D (PCHT 2D; OMIM #613811), a neurodegenerative condition characterized by progressive cerebrocerebellar atrophy, microcephaly, and epileptic encephalopathy. The identification of biallelic pathogenic variants in this family – one of which was a synonymous change not identified by prior clinical testing – not only ended the diagnostic odyssey for this family, but also highlights the contribution of occult pathogenic variants that may not be recognized by standard genetic testing methodologies.

CASE PRESENTATION

The 16-year-old female proband and 9-year-old male sibling described in this report were both delivered by spontaneous vaginal delivery with an unremarkable prenatal history. Hypotonia was noted shortly after birth. Their neonatal periods were marked by feeding issues and significant developmental delays by six months of age. The family history was considered non-contributory (Figure 1A). Their subsequent clinical course was nearly identical with dramatic reduction in head circumference, seizures associated with fever which later progressed into clusters of focal seizures. The proband's initial brain magnetic resonance imaging (MRI) indicated vasogenic edema of the frontal lobes probably secondary to venous sinus thrombosis. A follow up MRI indicated improvement. The initial diagnosis for the female proband was a static encephalopathy. However, the sibling was born with a similar phenotype with severe hypotonia, epilepsy and developmental delay.

At 16 years of age and 9 years of age the proband and male sibling respectively are described as having static encephalopathy, hypotonia, epilepsy, and intellectual disability (Table 1). Per clinician notes, both their facies are hypotonic with tented mouth, but otherwise exhibit no specific dysmorphic features. They are both non-verbal and non-ambulatory, with progressive chronic respiratory insufficiency, eventually requiring a tracheostomy and are currently ventilator dependent. They also have intractable epilepsy. See Supplemental Information for detailed clinical information.

Mitochondrial studies on the proband's muscle biopsy revealed reduction in mitochondrial complex I and II activity and an increase number of type 1 fibers in the muscle. Both siblings had extensive molecular testing including karyotype, oligonucleotide chromosomal microarray, Fragile X, and several single gene tests (see Supplemental Information), all of which was nondiagnostic. Following discussion of the case with the steering committee, the proband, sibling, mother, and father were enrolled in an IRB-approved research study (IRB11-00215) for genome sequencing.

TECHNICAL ANALYSIS AND METHODS

Genome Sequencing and Analysis

Following consent and enrollment, genomic DNA from the peripheral blood of proband, sibling and parents was extracted. Paired-end genome sequencing libraries were constructed using NEBNext Ultra II FS DNA Library Prep Kit (New England BioLabs) according to manufacturer's protocols. Whole-genome sequencing was performed using an Illumina NovaSeq6000 instrument according to manufacturer's protocols. Reads were mapped to the GRCh38 reference sequence and secondary data analysis was performed using our in-house Churchill pipeline (Kelly et al. 2015), which implements the GATK "best practices" workflow for alignment, variant discovery and genotyping. We generated ~136 Gbp per sample on average. Mapping the sequence reads to the GRCh38 reference sequence yielded ~41x haploid coverage per sample (range 33x to 53x; see Supplemental Table 1 for detailed metrics). Variants were joint-called in all family members using GATK 4.0.5.1, and the resulting VCF file was annotated with gene, transcript, function class, damaging scores, and population allele

frequencies using VarHouse, an in-house annotation tool built around the SNPeff annotation tool (Cingolani et al. 2012).

Variant annotation and prioritization were performed as previously described (Koboldt et al. 2018). Multi-sample variant calling with GATK HaplotypeCaller yielded 6,753,946 small sequence variants (SNVs and indels) in the family quad. Of these, 5,095 were in or near coding/UTR and splice site regions in the proband's sample and 2,414 were shared by the proband and sibling. After removing common variants (MAF>0.001 in the gnomAD and ExAC database), all splice site, frameshift, and nonsense variants, as well as missense variants predicted to be damaging by SIFT (score<0.05), Polyphen (score>0.453), GERP (score>2.0), or CADD (Phred score>15) were selected for further analysis. Given the family history, rare variants that segregated with disease under a recessive inheritance model (i.e., homozygous or compound-heterozygous) were prioritized and closely scrutinized. Out of 4 candidate genes (all in compound-heterozygous state in the siblings) that remained following our filtering, 2 were, at the time, not associated with human disease according to the OMIM database. The other two included *SEPSECS* (associated with autosomal recessive Pontocerebellar hypoplasia type 2D) and *DNAH17* (associated with autosomal recessive spermatogenic failure). As the latter was a poor phenotypic match for our proband and sibling's presentation, that left *SEPSECS* (Figure 1D) as the only candidate gene consistent with the expected inheritance model and matched the clinical presentation.

Variant Interpretation

The maternally inherited variant (also identified in a prior clinical WES on proband; see Supplemental information) is a novel SNV in the initiator codon (NM_016955.4:c.1A>T, evidence PVS1) that is absent from public databases (0 in gnomAD despite good coverage; evidence PM2). While this exact nucleotide change has not been previously reported, Zhu X. et al. (Zhu et al. 2015) identified another *SEPSECS* variant which affected the same codon (p.M1?) via a different nucleotide change (ENST00000382103.2; NM_016955.4: c.1A>G; evidence PS1). This change is predicted to result in protein translation start loss and is associated with a diagnosis of pontocerebellar hypoplasia type 2D in the proband (Zhu et al. 2015). The paternally inherited variant in *SEPSECS* (NM_016955.4: c.846G>A) is a rare SNV according to the gnomAD database (max population frequency 0.00019; evidence PM2). Although predicted to be a synonymous change, this variant was shown experimentally to disrupt mRNA splicing (evidence PS3) using affected proband's blood RNASeq data in a recent study by the Undiagnosed Diseases Network (Lee et al. 2020) and reported as likely pathogenic to the ClinVar database (evidence PP5; ClinVar ID-SCV000746576.2). Briefly, Lee et al. demonstrated that the synonymous variant (located on exon 7 of *SEPSECS* gene product) which was 42 bp downstream of a splice acceptor site caused exon 7 skipping in nearly all of the *SEPSECS* transcripts in proband and mother. Though there was an indication of exon 7 skipping in unrelated control samples, this occurred in significantly lesser number of transcripts suggesting that this exon is prone to skipping and the synonymous change greatly increased the chances of this event (Lee et al. 2020). Though the mechanisms underlying this altered splicing event is unclear, one possible explanation for this skipping could be that this synonymous change maybe altering an exonic splice enhancer or silencer within the exon. Ramser et al. (Ramser et al. 2005) have previously reported a synonymous change in the *ATP6AP2* gene, leading to exon skipping via the alteration of a putative exonic splice enhancer. Further, such exon skipping events are well-established disease-causing mechanisms, as seen in spinal muscular atrophy and other

conditions (Lorson et al. 1999; Anna and Monika 2018). Though this protein change is not novel, ours is the first study that reports the occurrence of this variant in combination with a different *SEPSECS* variant *in trans*. Both variants were compound heterozygous in the two affected children, consistent with recessive inheritance (evidence PP1). Taken together, under ACMG guidelines, the maternally inherited variant was classified ‘pathogenic’ while the paternally inherited variant was classified ‘likely pathogenic’ (Table 2).

These genomic findings were discussed with the referring clinician, who concluded that the clinical presentation of the siblings was a good match to the phenotype of pontocerebellar hypoplasia type 2D associated with *SEPSECS* gene variants. The results were shared with the family, who declined segregation testing for the unaffected siblings.

SUMMARY

Pontocerebellar hypoplasia (PCH; OMIM #607596) is a group of heterogenous and often lethal autosomal recessive neurodegenerative disorders that affect the growth and function of the pons (a major part of the brainstem) and cerebellum. Since its initial classification into type 1 and 2 PCH based on accompanying clinical phenotypes; (Barth 1993), there are currently nine different subtypes with varying clinical presentations and genetic causes. Specifically, pontocerebellar hypoplasia type 2 (PCH2) involves a lack of voluntary motor development. Individuals with PCH2 present with impaired vision, dysphagia, severe impairment of cognitive and gross motor function, and seizures of varying severity (Namavar et al. 2011). There are currently variants in six unique genes are known to cause PCH2, resulting in a sub-classification of A through F (van Dijk et al. 2018). The first report of PCH2D (OMIM #613811) was found in a cohort of Sephardic Jews who presented with intellectual disability, progressive microcephaly, spasticity, and progressive cerebellar-cerebral atrophy (Ben-Zeev et al. 2003) and was later linked to *SEPSECS* gene (Agamy et al. 2010). Pontocerebellar hypoplasia type 2D is characterized by progressive microcephaly, cerebellar atrophy, early onset epileptic encephalopathy, and optic nerve atrophy (van Dijk et al. 2018) complicated by heterogeneity in symptoms and severity. Our patients’ presentation was consistent with the classical phenotypic spectrum comprising progressive microcephaly, cerebellar atrophy (Figure 1B, Supplemental Figures 1&2), vision loss, epilepsy, and dysphagia.

The identification of biallelic pathogenic *SEPSECS* variants (one of which was a synonymous change) in the siblings led to a critical diagnostic breakthrough for the family and demonstrates the contribution of occult splicing variants to monogenic disorders.

TABLES

Table 1. Patient clinical characteristics using HPO terms where “+” indicates presence, “-” indicates absence, “unk” indicates unknown.

HPO ID	Phenotype	Proband, 16yo female	Sibling, 9yo male
HP:0001290	Generalized hypotonia	+	+
HP:0002194	Delayed gross motor development	+	+
HP:0001263	Global developmental delay	+	+
HP:0002187	Intellectual disability; profound	+	+
HP:0012736	Profound global developmental delay	+	+

HP:0001250	Seizures	+	+
HP:0002650	Scoliosis	+	+
HP:0000505	Visual impairment	+	+
HP:0002579	Gastrointestinal dysmotility	+	+
HP:0002093	Respiratory insufficiency	+	+
HP:0011922	Abnormal activity of mitochondrial respiratory chain	+	unk
HP:0008347	Decreased activity of mitochondrial complex IV	+	unk
HP:0000252	Microcephaly	+	+
HP:0008936	Muscular hypotonia of the trunk	+	+
HP:008872	Feeding difficulties in infancy	+	+

Table 2. Genomic findings and variant interpretation. Genomic coordinates reflect build GRCh38.

Gene	Variant	HGVS	Proband	Mother	Father	Sibling	Interpretation
<i>SEPSECS</i>	chr4: 25160369 T/A	NM_016955.4: c.1A>T	Het	Het	Ref	Het	P (PVS1, PS1, PM2, PP1)
<i>SEPSECS</i>	chr4: 25145092 C/T	NM_016955.4: c.846G>A	Het	Ref	Het	Het	LP (PS3, PM2, PP1, PP5)

FIGURES

Figure 1. Whole genome sequencing of family quad reveals biallelic variants in *SEPSECS* gene in proband and sibling. A. Pedigree of the affected siblings. B. Diffuse cerebral and

cerebellar cortical and white matter volume loss observed in male sibling's MRI. C. Lollipop plots that map all known pathogenic/likely pathogenic SEPSECS variants found in the ClinVar database as of January 1, 2021 as well as novel variants found in this study (black dots) on the exons encoding SEPSECS protein. D. Confirmatory Sanger sequencing traces (reverse primer) demonstrating the biallelic SEPSECS variants in both siblings while unaffected parents were both carriers for one of the SEPSECS variant alleles. Highlighted bases indicate the single nucleotide variation (SNV) in respective family members.

REFERENCES

- Agamy, O., B. Ben Zeev, D. Lev, B. Marcus, D. Fine, D. Su, G. Narkis, R. Ofir, C. Hoffmann, E. Leshinsky-Silver, H. Flusser, S. Sivan, D. Söll, T. Lerman-Sagie, and O. S. Birk. 2010. 'Mutations disrupting selenocysteine formation cause progressive cerebello-cerebral atrophy', *Am J Hum Genet*, 87: 538-44.
- Anna, A., and G. Monika. 2018. 'Splicing mutations in human genetic disorders: examples, detection, and confirmation', *J Appl Genet*, 59: 253-68.
- Barth, P. G. 1993. 'Pontocerebellar hypoplasias. An overview of a group of inherited neurodegenerative disorders with fetal onset', *Brain Dev*, 15: 411-22.
- Ben-Zeev, B., C. Hoffman, D. Lev, N. Watemberg, G. Malinger, N. Brand, and T. Lerman-Sagie. 2003. 'Progressive cerebellocerebral atrophy: a new syndrome with microcephaly, mental retardation, and spastic quadriplegia', *J Med Genet*, 40: e96.
- Cingolani, P., A. Platts, L. Wang le, M. Coon, T. Nguyen, L. Wang, S. J. Land, X. Lu, and D. M. Ruden. 2012. 'A program for annotating and predicting the effects of single nucleotide polymorphisms, SnpEff: SNPs in the genome of *Drosophila melanogaster* strain w1118; iso-2; iso-3', *Fly (Austin)*, 6: 80-92.
- Kelly, B. J., J. R. Fitch, Y. Hu, D. J. Corsmeier, H. Zhong, A. N. Wetzel, R. D. Nordquist, D. L. Newsom, and P. White. 2015. 'Churchill: an ultra-fast, deterministic, highly scalable and balanced parallelization strategy for the discovery of human genetic variation in clinical and population-scale genomics', *Genome Biol*, 16: 6.
- Koboldt, D. C., T. Mihalic Mosher, B. J. Kelly, E. Sites, D. Bartholomew, S. E. Hickey, K. McBride, R. K. Wilson, and P. White. 2018. 'A de novo nonsense mutation in ASXL3 shared by siblings with Bainbridge-Ropers syndrome', *Cold Spring Harb Mol Case Stud*, 4.
- Lee, H., A. Y. Huang, L. K. Wang, A. J. Yoon, G. Renteria, A. Eskin, R. H. Signer, N. Dorrani, S. Nieves-Rodriguez, J. Wan, E. D. Douine, J. D. Woods, E. C. Dell'Angelica, B. L. Fogel, M. G. Martin, M. J. Butte, N. H. Parker, R. T. Wang, P. B. Shieh, D. A. Wong, N. Gallant, K. E. Singh, Y. J. Tavyev Asher, J. S. Sinsheimer, D. Krakow, S. K. Loo, P. Allard, J. C. Papp, C. G. S. Palmer, J. A. Martinez-Agosto, S. F. Nelson, and Undiagnosed Diseases Network. 2020. 'Diagnostic utility of transcriptome sequencing for rare Mendelian diseases', *Genet Med*, 22: 490-99.
- Lorson, C. L., E. Hahnen, E. J. Androphy, and B. Wirth. 1999. 'A single nucleotide in the SMN gene regulates splicing and is responsible for spinal muscular atrophy', *Proc Natl Acad Sci U S A*, 96: 6307-11.
- Namavar, Y., P. G. Barth, B. T. Poll-The, and F. Baas. 2011. 'Classification, diagnosis and potential mechanisms in pontocerebellar hypoplasia', *Orphanet J Rare Dis*, 6: 50.

- Ramser, J., F. E. Abidi, C. A. Burckle, C. Lenski, H. Toriello, G. Wen, H. A. Lubs, S. Engert, R. E. Stevenson, A. Meindl, C. E. Schwartz, and G. Nguyen. 2005. 'A unique exonic splice enhancer mutation in a family with X-linked mental retardation and epilepsy points to a novel role of the renin receptor', *Hum Mol Genet*, 14: 1019-27.
- van Dijk, T., F. Baas, P. G. Barth, and B. T. Poll-The. 2018. 'What's new in pontocerebellar hypoplasia? An update on genes and subtypes', *Orphanet J Rare Dis*, 13: 92.
- Zhu, X., S. Petrovski, P. Xie, E. K. Ruzzo, Y. F. Lu, K. M. McSweeney, B. Ben-Zeev, A. Nissenkorn, Y. Anikster, D. Oz-Levi, R. S. Dhindsa, Y. Hitomi, K. Schoch, R. C. Spillmann, G. Heimer, D. Marek-Yagel, M. Tzadok, Y. Han, G. Worley, J. Goldstein, Y. H. Jiang, D. Lancet, E. Pras, V. Shashi, D. McHale, A. C. Need, and D. B. Goldstein. 2015. 'Whole-exome sequencing in undiagnosed genetic diseases: interpreting 119 trios', *Genet Med*, 17: 774-81.

ADDITIONAL INFORMATION

Ethics Statement

Written informed consent was obtained for all participants in this study under a research protocol approved by the Institutional Review Board at Nationwide Children's Hospital (IRB11-00215 Study: Using Genome Sequencing to Identify Causes of Rare Birth Defects and Rare Disorders).

Data Deposition

The SEPSECS variants and our interpretation have been submitted to the ClinVar database (Accession IDs: SCV001960820.1 (NM_016955.4:c.1A>T) and SCV001960821.1 (NM_016955.4:c.846G>A)).

Author Contributions

All authors contributed to scientific discussion, variant interpretation, and manuscript review.

Acknowledgments

We thank the family for participation in this research.

Funding

This work was supported by The Research Institute at Nationwide Children's Hospital.

Biallelic SEPSECS variants in two siblings with pontocerebellar hypoplasia type 2D underscore the relevance of splice-disrupting synonymous variants in disease.

Swetha Ramadesikan, Scott E Hickey, Emily De Los Reyes, et al.

Cold Spring Harb Mol Case Stud published online January 28, 2022
Access the most recent version at doi:[10.1101/mcs.a006165](https://doi.org/10.1101/mcs.a006165)

Supplementary Material

<http://molecularcasestudies.cshlp.org/content/suppl/2022/02/04/mcs.a006165.DC1>

Published online January 28, 2022 in advance of the full issue.

Accepted Manuscript

Peer-reviewed and accepted for publication but not copyedited or typeset; accepted manuscript is likely to differ from the final, published version. Published online January 28, 2022 in advance of the full issue.

Creative Commons License

This article is distributed under the terms of the <http://creativecommons.org/licenses/by-nc/4.0/>, which permits reuse and redistribution, except for commercial purposes, provided that the original author and source are credited.

Email Alerting Service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#).